

学校编码: 10384

密级_____

学号: 33320131151737

廈門大學

硕士学位论文

同安湾下潭尾人工红树林湿地大型底栖动物群落恢复研究

Study on restoration of benthic macrofaunal community in
Xiatanwei artificial mangrove wetlands, Tong'an Bay

颜露露

指导教师姓名: 蔡立哲 教授

专业名称: 环境科学

论文提交日期: 2016 年 4 月

论文答辩时间: 2016 年 5 月

2016 年 5 月

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下,独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果,均在文中以适当方式明确标明,并符合法律规范和《厦门大学研究生学术活动规范(试行)》。

另外,该学位论文为()课题(组)的研究成果,获得()课题(组)经费或实验室的资助,在()实验室完成。(请在以上括号内填写课题或课题组负责人或实验室名称,未有此项声明内容的,可以不作特别声明。)

声明人(签名):

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文，并向主管部门或其指定机构送交学位论文（包括纸质版和电子版），允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索，将学位论文的标题和摘要汇编出版，采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于：

（ ） 1. 经厦门大学保密委员会审查核定的保密学位论文，
于 年 月 日解密，解密后适用上述授权。

（ ） 2. 不保密，适用上述授权。

（请在以上相应括号内打“√”或填上相应内容。保密学位论文应是已经厦门大学保密委员会审定过的学位论文，未经厦门大学保密委员会审定的学位论文均为公开学位论文。此声明栏不填写的，默认为公开学位论文，均适用上述授权。）

声明人（签名）：

年 月 日

目 录

摘要.....	I
Abstract.....	III
第一章 前言.....	6
1.1 国内外潮间带大型底栖动物研究进展.....	6
1.1.1 国内潮间带大型底栖动物研究进展.....	7
1.1.2 国外潮间带大型底栖动物群落研究进展.....	8
1.2 国内红树林生态系统大型底栖动物群落研究进展.....	11
1.2.1 国内红树林生态系统及其生态恢复研究进展.....	11
1.2.2 国内红树林生态系统大型底栖动物群落研究进展.....	12
1.3 同安湾底栖动物群落生态研究进展.....	13
1.3.1 同安湾海域概况.....	13
1.3.2 同安湾潮间带和红树林区底栖动物群落生态研究进展.....	14
1.3.3 同安湾潮下带底栖动物群落生态研究进展.....	15
1.4 大型底栖动物功能群研究概况.....	17
1.5 研究意义和技术路线.....	18
第二章 材料与方法.....	21
2.1 同安湾下潭尾人工红树林湿地概况及大型底栖动物取样站设置.....	21
2.1.1 同安湾下潭尾人工红树林湿地自然地理概况.....	21
2.1.2 同安湾下潭尾人工红树林湿地大型底栖动物取样站设置.....	21
2.2 人工红树林湿地大型底栖动物的采集、保存处理和数据统计方法.....	22
2.3 大型底栖动物群落结构分析方法.....	23
2.3.1 群落结构分析.....	23
2.3.2 大型底栖动物群落物种多样性单变量分析.....	24
2.3.3 单变量双因素方差分析 (Two-way ANOVA)	25
2.4 水质和沉积物环境因子的测定.....	25
2.5 大型底栖动物功能群划分方法.....	26

2.6 大型底栖动物群落健康评价.....	27
2.6.1 丰度-生物量比较法.....	27
2.6.2 大型底栖动物污染指数法.....	28
2.6.3 海洋生物指数和多变量海洋生物指数.....	28
第三章 同安湾下潭尾人工红树林湿地大型底栖动物群落.....	31
3.1 同安湾下潭尾人工红树林湿地的大型底栖动物种类组成.....	31
3.2 同安湾下潭尾人工红树林湿地的大型底栖动物栖息密度.....	32
3.3 同安湾下潭尾人工红树林湿地大型底栖动物生物量的时空分布.....	34
3.4 讨论.....	36
3.4.1 同安湾下潭尾人工红树林湿地的大型底栖动物群落比较.....	36
3.4.2 与同安湾其他红树林区大型底栖动物群落的比较.....	37
第四章 同安湾下潭尾人工红树林湿地大型底栖动物功能群.....	39
4.1 同安湾下潭尾人工红树林湿地大型底栖动物功能群组成.....	39
4.1.1 同安湾下潭尾人工红树林湿地大型底栖动物功能群的物种组成	39
4.1.2 同安湾下潭尾人工红树林湿地大型底栖动物功能群的栖息密度	40
4.1.3 同安湾下潭尾人工红树林湿地大型底栖动物功能群的生物量.....	42
4.2 下潭尾人工红树林湿地大型底栖动物功能群的多样性指数.....	43
4.2.1 同安湾下潭尾人工红树林湿地大型底栖动物功能群的种类多样性 指数 H'	43
4.2.2 同安湾下潭尾人工红树林湿地大型底栖动物功能群的种类丰富度 指数 d	44
4.2.3 同安湾下潭尾人工红树林湿地大型底栖动物功能群的种类均匀度 指数 J	44
4.3 讨论.....	47
4.3.1 生境差异是影响同安湾下潭尾人工红树林湿地大型底栖动物功能 群结构的主要因素.....	47
4.3.2 某些大型底栖动物食性归属难以确定影响功能群的划分.....	47
第五章 下潭尾人工红树林区大型底栖动物群落的边缘效应.....	49

5.1 人工红树林区林中与林缘大型底栖动物群落比较.....	49
5.1.1 人工红树林区林中与林缘大型底栖动物的种类组成.....	49
5.1.2 人工红树林区林中与林缘大型底栖动物的栖息密度组成.....	50
5.1.3 人工红树林区林中与林缘大型底栖动物的生物量组成.....	51
5.2 人工红树林区林内与林缘大型底栖动物的多样性指数比较.....	52
5.3 人工红树林区林内与林缘大型底栖动物的次级生产力比较.....	53
5.3.1 大型底栖动物次级生产力的计算.....	53
5.3.2 人工红树林区林内与林缘大型底栖动物的次级生产力.....	54
5.4 讨论.....	55
第六章 下潭尾人工红树林湿地大型底栖动物群落健康评价.....	57
6.1 群落物种生物多样性单变量分析.....	57
6.1.1 下潭尾人工红树林湿地大型底栖动物 Shannon-Wiener 种类多样性指数.....	57
6.1.2 同安湾下潭尾人工红树林湿地大型底栖动物群落的种类丰富度指数.....	58
6.1.3 同安湾下潭尾人工红树林湿地大型底栖动物群落的均匀度指数.....	59
6.2 同安湾下潭尾人工红树林湿地大型底栖动物 MPI 指数.....	59
6.3 同安湾下潭尾人工红树林湿地大型底栖动物的 AMBI 值.....	60
6.4 同安湾下潭尾人工红树林湿地大型底栖动物的 M-AMBI 值.....	60
6.5 下潭尾人工红树林湿地大型底栖动物的丰度-生物量曲线.....	61
6.6 同安湾下潭尾人工红树林湿地大型底栖动物群落的相似性分析.....	64
6.7 讨论.....	66
6.7.1 同安湾下潭尾人工红树林湿地大型底栖动物群落健康评价.....	66
6.7.2 大型底栖动物群落健康评价方法的比较.....	66
第七章 同安湾下潭尾人工红树林湿地大型底栖动物与环境变量关系分析.....	68
7.1 同安湾下潭尾人工红树林湿地理化因子和沉积物环境状况.....	68
7.1.1 同安湾下潭尾人工红树林湿地沉积物的温度和盐度.....	68

7.1.2 同安湾下潭尾人工红树林湿地沉积物环境状况.....	69
7.1.3 同安湾下潭尾人工红树林湿地沉积物的粒径组成.....	70
7.2 同安湾下潭尾人工红树林湿地大型底栖动物群落与环境因子的相关分析.....	71
7.2.1 同安湾下潭尾人工红树林湿地大型底栖动物群落与水质环境因子的关系.....	71
7.2.2 同安湾下潭尾人工红树林湿地大型底栖动物群落与沉积物环境因子的相关分析.....	72
7.2.3 同安湾下潭尾人工红树林湿地大型底栖动物群落与沉积物粒径组成的相关分析.....	73
7.3 讨论.....	74
7.3.1 盐度对下潭尾人工红树林湿地大型底栖动物群落的影响.....	74
7.3.2 底质粒径对大型底栖动物群落的影响.....	75
第八章 论文小结.....	77
8.1 研究成果.....	77
8.2 创新点.....	78
8.3 不足之处.....	78
8.4 展望.....	79
参考文献.....	80
附表 1 同安湾人工红树林湿地大型底栖动物名录.....	92
参加的课题及发表的文章.....	98
致谢.....	99

Table of contents

Abstract(In Chinese).....	I
Abstract(In English).....	III
Chapter 1 Preface.....	6
1.1 Research progress on benthic macrofauna in domestic and foreign intertidal areas.....	6
1.1.1 Research progress on benthic macrofauna in domestic intertidal areas.....	7
1.1.2 Research progress on benthic macrofauna in foreign intertidal areas.....	8
1.2 Research progress on benthic macrofaunal community in domestic mangrove ecosystem.....	11
1.2.1 Research progress on domestic mangrove ecosystem and its ecological restoration.....	11
1.2.2 Research progress on benthic macrofaunal community in domestic mangrove ecosystem.....	12
1.3 Research progress on zoobenthic community ecology in Tong'an Bay.....	13
1.3.1 Geography overview of Tong'an Bay.....	13
1.3.2 Research progress on zoobenthic community ecology in intertidal and mangrove areas in Tong'an Bay.....	14
1.3.3 Research progress on zoobenthic community ecology in subtidal zones in Tong'an Bay.....	15
1.4 Research overview on functional groups of benthic macrofaunal	17
1.5 Significance and technological route in this study.....	18
Chapter 2 Materials and methods.....	21
2.1 General situation of Xiatanwei artificial mangrove wetlands in Tong'an Bay and sampling stations setting.....	21
2.1.1 Physical geography situation of Xiatanwei artificial mangrove wetlands.....	21
2.1.2 Sampling stations setting of Xiatanwei artificial mangrove wetlands.....	21
2.2 Sampling, preservation and processing statistical method of benthic macrofauna in artificial mangrove wetlands.....	22
2.3 Methods of benthic macrofaunal community statistical analysis.....	23
2.3.1 Analysis of benthic macrofaunal community structure.....	23
2.3.2 Univariate variable analysis of species diversity.....	24
2.3.3 Two-way analysis of variance (Two-way ANOVA).....	25
2.4 Determination of seawater and sediment environmental factors.....	25
2.5 Division of benthic macrofaunal functional groups.....	26
2.6 Health assessment methods of benthic macrofaunal community	27
2.6.1 The curve of abundance and biomass comparison.....	27
2.6.2 Macrozoobenthos Pollution Index (MPI).....	28
2.6.3 AZTI'S Marine Biotic Index (AMBI) and Multivariate AZTI'S Marine Biotic Index (M-AMBI).....	28

Chapter 3 Benthic macrofaunal community in Xiatanwei artificial

mangrove wetlands in Tong'an Bay.....	31
3.1 Species composition of benthic macrofaunal community	31
3.2 Spatial-temporal distribution of benthic macrofaunal density	32
3.3 Spatial-temporal distribution of benthic macrofaunal biomass	34
3.4 Discussion.....	36
3.4.1 Comparison of benthic macrofaunal community in Xiatanwei artificial mangrove wetlands in Tong'an Bay.....	36
3.4.2 Comparison of benthic macrofaunal community among different mangrove areas.....	37

Chapter 4 Benthic macrofaunal functional groups in Xiatanwei artificial

mangrove wetlands in Tong'an Bay.....	39
4.1 Composition of benthic macrofaunal functional groups in Xiatanwei artificial mangrove wetlands in Tong'an Bay.....	39
4.1.1 Species composition of benthic macrofaunal functional groups.....	39
4.1.2 Density composition of benthic macrofaunal functional groups	40
4.1.3 Biomass composition of benthic macrofaunal functional groups	42
4.2 Diversity indices of benthic macrofaunal functional groups	43
4.2.1 Shannon-Wiener diversity index(H')	43
4.2.2 Margalef species abundance index(d)	44
4.2.3 Pielous species evenness index(J)	44
4.3 Discussion.....	47
4.3.1 Habitat differences influenced the structure of benthic macrofaunal functional groups.....	47
4.3.2 Feeding type of some species was difficult to determine affected the result of functional groups research.....	47

Chapter 5 Edge effect of benthic macrofaunal community in Xiatanwei

artificial mangrove wetlands in Tong'an Bay.....	49
5.1 Comparison of benthic macrofaunal community between the forest interior and edges of artificial mangrove wetlands.....	49
5.1.1 Species composition of benthic macrofaunal community	49
5.1.2 Density composition of benthic macrofaunal community	50
5.1.3 Biomass composition of benthic macrofaunal community.....	51
5.2 Comparison of benthic macrofaunal diversity indices between the forest interior and edges of artificial mangrove wetlands.....	52
5.3 Comparison of benthic macrofaunal secondary productivity between the forest interior and edges of artificial mangrove wetlands.....	53
5.3.1 Calculation method of secondary productivity.....	53

5.3.2	Comparison of benthic macrofaunal secondary productivity	54
5.4	Discussion.....	55
Chapter 6 Health assessment of benthic macrofaunal community in		
	Xiatanwei artificial mangrove wetlands in Tong'an Bay.....	57
6.1	Univariate analysis of species biodiversity.....	57
6.1.1	Shannon-Wiener diversity index(H') of benthic macrofaunal community.....	57
6.1.2	Margalef species abundance index(d) of benthic macrofaunal community.....	58
6.1.3	Pielous species evenness index(J) of benthic macrofauna community.....	59
6.2	Macrozoobenthos Pollution Index (MPI).....	59
6.3	AZTI'S Marine Biotic Index	60
6.4	Multivariate AZTI'S Marine Biotic Index.....	60
6.5	The curve of abundance and biomass comparison.....	61
6.6	Similarity analysis of benthic macrofaunal community in Xiatanwei artificial mangrove wetlands in Tong'an Bay.....	64
6.7	Discussion.....	66
6.7.1	Health assessment of benthic macrofaunal community in Xiatanwei artificial mangrove wetlands in Tong'an Bay.....	66
6.7.2	Comparison of benthic macrofaunal community health assessment methods.....	66
Chapter 7 The relationship between benthic macrofaunal community		
	and environment factors.....	68
7.1	Physical-chemical factors and environmental status of sediment in Xiatanwei artificial mangrove wetlands in Tong'an Bay.....	68
7.1.1	Temperature and Salinity.....	68
7.1.2	Environmental status of sediment	69
7.1.3	Sedimental grain size.....	70
7.2	Pearson correlation between benthic macrofaunal community and environment factors in Xiatanwei artificial mangrove wetlands in Tong'an Bay.....	71
7.2.1	Pearson correlation between benthic macrofaunal community and seawater environment factors.....	71
7.2.2	Pearson correlation between benthic macrofaunal community and sediment environment factors.....	72
7.2.3	Pearson correlation between benthic macrofaunal community and sedimental grain size	73
7.3	Discussion.....	74
7.3.1	Effects of salinity on benthic macrofaunal community.....	74
7.3.2	Effects of sedimental grain size on benthic macrofaunal community.....	75
Chapter 8 Summary.....		
8.1	Study results.....	77
8.2	Innovation.....	78

8.3 Insufficiency.....	78
8.4 Prospection.....	79
Reference.....	80
Appendix.....	92
Participation issues and articles published.....	98
Acknowledgements.....	99

厦门大学博硕士论文摘要库

摘要

红树林生态系统处于海洋与陆地的动态交界面,是我国三大海岸带生态系统之一。底栖动物(包括大型、小型和微型底栖动物)是红树林生态系统的重要组成部分,同时也是该生态系统内物质循环、能量流动中积极的消费者和转移者。关于同安湾潮间带、潮下带和红树林湿地大型底栖动物群落生态的研究已有报道,但关于同安湾人工红树林湿地大型底栖动物群落恢复、演替和功能群方面的研究稀少,且关于红树林群落边缘大型底栖动物的群落效应的研究未见报道。

本研究选取同安湾下潭尾人工红树林湿地作为研究区域,比较了不同树龄红树林湿地大型底栖动物群落的时空变化、功能群组成、生物指数和边缘效应,分析了大型底栖动物群落参数与人工红树林湿地环境变量之间的关系,对比了生物指数、丰度生物量比较法评价人工红树林湿地生态环境现状的结果。研究结果可为红树林生态系统的恢复和管理、大型底栖动物资源的开发利用提供科学依据。主要研究成果如下:

1. 2014 年 4 月至 2015 年 1 月在同安湾下潭尾人工红树林湿地共获得大型底栖动物 93 种,其中定性采集到 40 种,定量采集到 65 种。定量和定性采集获得的共有种 12 种,仅占定量采集物种数的 18.46%、定性采集物种数的 30.00%。定量采集的大型底栖动物物种数以多毛类和甲壳类居多,均有 24 种;定性采集的大型底栖动物物种数以甲壳类为主,有 26 种。

2. 同安湾下潭尾人工红树林湿地大型底栖动物的年平均栖息密度为 $695.4 \text{ ind.} \cdot \text{m}^{-2}$ 。从树龄分布看,2 a 树龄红树林区大型底栖动物栖息密度最高,其次分别是光滩、1 a 林、9 a 林;从季节分布看,春季最高,其次分别是冬季、夏季和秋季。年平均生物量为 $24.07 \text{ g} \cdot \text{m}^{-2}$ 。从树龄上看,光滩大型底栖动物年平均生物量最高,其次分别是 2 a 林、9 a 林、1 a 林;从季节分布看,春季最高,其次分别是秋季、夏季和冬季。

3. 同安湾下潭尾人工红树林湿地定量采集的大型底栖动物功能群以杂食者为主要类型,达到 21 种;浮游生物食者物种数最少,只有 6 种。生境差异是影响该区域大型底栖动物功能群结构的主要因素。

4. 根据大型底栖动物群落评价人工红树林区边缘效应的强弱,发现林缘大

型底栖动物的物种数、栖息密度、物种丰富度和次级生产力均高于林内，证明边缘对该林区的大型底栖动物群落产生了正效应。

5. 运用生物多样性指数法、大型底栖动物污染指数 MPI 法、AMBI(海洋生物指数)和 M-AMBI(多变量海洋生物指数)指数法以及丰度-生物量比较 (ABC) 曲线法对同安湾下潭尾人工红树林湿地大型底栖动物群落进行环境健康评价, 结果显示 2 a 红树林区处于轻度扰动状态, 其他 3 种树龄红树林区处于无扰动和轻度扰动状态; 同时 ABC 曲线法对 1 a 红树林区、2 a 红树林区以及 9 a 红树林区的评价结果与其他 3 类评价方法存在偏差, 这可能与 ABC 曲线法在应用时存在局限性有关。

6. 分析同安湾下潭尾人工红树林湿地大型底栖动物群落与环境变量之间的关系, 发现大型底栖动物群落的季节变化与海水盐度之间存在显著或极显著负相关关系, 同时沉积物底质差异影响大型底栖动物物种数、栖息密度和生物量在不同龄红树林区的分布。

关键词: 大型底栖动物; 人工红树林湿地; 群落恢复; 同安湾

Abstract

Mangrove system is one of China's three major coastal ecosystems, usually situating in the dynamic interface between the marine and terrestrial ecosystems. Macrofauna is an important component of mangrove ecosystem, as well as a positive consumer and metastasis group in the material cycle and energy flow process. There are many researches about benthic macrofaunal community in the intertidal zones, subtidal zones and mangrove wetlands in Tong'an Bay, but researches on community restoration, succession and functional groups of benthic macrofauna in artificial mangrove wetlands in Tong'an Bay are relatively scarce.

This study selecting the artificial mangrove wetlands in Tong'an Bay as the study area, comparing the temporal and spatial variation of macrofaunal community among mudflat, 1-year mangrove forest, 2-year mangrove forest and 9-year mangrove forest. The relationship between environment variable and the macrofaunal community in different biotopes was analyzed. Simultaneously, this study comparing the biological indices and functional group composition of macrofaunal community in the artificial mangrove of different age, then contrasting the results of different evaluation methods. It was the first time to use the benthic macrofaunal community to explore the strength of edge effect in artificial mangrove forest. The results from the present study can provide a scientific basis for mangrove restoration and management, as well as the exploitation and utilization of macrofauna resources. The main results are as follows:

1. Ninety-three species of benthic macrofauna were obtained in the artificial mangrove wetlands of Tong'an Bay, including 40 species were obtained from the qualitative collection and 65 species were obtained from the quantitative collection. Collection of qualitative and quantitative shared a total of 12 common species, accounting for 18.46 percent of the species number of quantitative collection, and 30 percent of the species number of qualitative collection. Polychaetes and crustaceans had a majority in the quantitative collection of benthic macrofauna, both of them including 24 species. Crustaceans that amounted to 26 species was the dominant

group in qualitative sampling.

2. The annual average density of benthic macrofauna in the study area was 695.4 ind. \cdot m⁻², in which the 2-year mangrove forest had the highest density, followed by mudflat, 1-year mangrove forest and 9-year mangrove forest. Among the seasonal distribution, spring had the highest density, followed by winter, summer and autumn. The annual average biomass of benthic macrofauna was 24.07 g \cdot m⁻², the mudflat habitat shared the highest biomass, in which mudflat had the highest biomass, followed by 2-year mangrove forest, 9-year mangrove forest and 1-year mangrove forest. Among the seasonal distribution, the highest biomass appeared in spring, followed by autumn, summer and winter.

3. Omnivores group was the main benthic macrofaunal functional group in Tong'an Bay artificial mangrove wetlands, and planktonphagous group had the least number of species, only 6 species. Habitat diversity was a major factor that impacted the benthic macrofaunal functional group structure.

4. The edge effect of mangrove area was first discussed according to benthic macrofaunal community. It was found that species number, density, species richness and secondary productivity of benthic macrofauna in the forest edge were higher than in the forest interior. This result proved that the edge of mangrove forest had a positive effect to benthic macrofaunal community.

5. Biological diversity index, Macrofaunal Pollution Index (MPI), AZTI'S Marine Biotic Index (AMBI), Multivariate AZTI'S Marine Biotic Index (M-AMBI), and Abundance Biomass Comparison curve (ABC method) were applied to evaluate the environmental health of benthic macrofaunal community in Xiatanwei artificial mangrove wetlands, Tong'an Bay. The results showed that 2-year mangrove forest was slightly disturbed, while the other three biotopes were in undisturbed or mildly disturbed state. Meanwhile the evaluation result of ABC curve in the study area showed a deviation with other methods, which may be related to the limitation of ABC curve.

6. The relationship between benthic macrofaunal community in the artificial mangrove wetlands and environment variables was analyzed, it showed that there was

significant and extremely significant negative correlation between the seasonal variation of benthic macrofaunal community and seawater salinity, while the deposit sediment condition affect the species number, density, biomass of benthic macrofauna in different mangrove areas.

Key Words: Benthic macrofauna, Artificial mangrove wetlands, Community restoration, Tong'an Bay

Degree papers are in the “[Xiamen University Electronic Theses and Dissertations Database](#)”.

Fulltexts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.